



Sustento del uso justo
de Materiales Protegidos
derechos de autor para
fines educativos



UCI

Universidad para la
Cooperación Internacional

UCI
Sustento del uso justo de materiales protegidos por
derechos de autor para fines educativos

El siguiente material ha sido reproducido, con fines estrictamente didácticos e ilustrativos de los temas en cuestión, se utilizan en el campus virtual de la Universidad para la Cooperación Internacional – UCI – para ser usados exclusivamente para la función docente y el estudio privado de los estudiantes pertenecientes a los programas académicos.

La UCI desea dejar constancia de su estricto respeto a las legislaciones relacionadas con la propiedad intelectual. Todo material digital disponible para un curso y sus estudiantes tiene fines educativos y de investigación. No media en el uso de estos materiales fines de lucro, se entiende como casos especiales para fines educativos a distancia y en lugares donde no atenta contra la normal explotación de la obra y no afecta los intereses legítimos de ningún actor.

La UCI hace un USO JUSTO del material, sustentado en las excepciones a las leyes de derechos de autor establecidas en las siguientes normativas:

- a- Legislación costarricense: Ley sobre Derechos de Autor y Derechos Conexos, No.6683 de 14 de octubre de 1982 - artículo 73, la Ley sobre Procedimientos de Observancia de los Derechos de Propiedad Intelectual, No. 8039 – artículo 58, permiten el copiado parcial de obras para la ilustración educativa.
- b- Legislación Mexicana; Ley Federal de Derechos de Autor; artículo 147.
- c- Legislación de Estados Unidos de América: En referencia al uso justo, menciona: "está consagrado en el artículo 106 de la ley de derecho de autor de los Estados Unidos (U.S, Copyright - Act) y establece un uso libre y gratuito de las obras para fines de crítica, comentarios y noticias, reportajes y docencia (lo que incluye la realización de copias para su uso en clase)."
- d- Legislación Canadiense: Ley de derechos de autor C-11– Referidos a Excepciones para Educación a Distancia.
- e- OMPI: En el marco de la legislación internacional, según la Organización Mundial de Propiedad Intelectual lo previsto por los tratados internacionales sobre esta materia. El artículo 10(2) del Convenio de Berna, permite a los países miembros establecer limitaciones o excepciones respecto a la posibilidad de utilizar lícitamente las obras literarias o artísticas a título de ilustración de la enseñanza, por medio de publicaciones, emisiones de radio o grabaciones sonoras o visuales.

Además y por indicación de la UCI, los estudiantes del campus virtual tienen el deber de cumplir con lo que establezca la legislación correspondiente en materia de derechos de autor, en su país de residencia.

Finalmente, reiteramos que en UCI no lucramos con las obras de terceros, somos estrictos con respecto al plagio, y no restringimos de ninguna manera el que nuestros estudiantes, académicos e investigadores accedan comercialmente o adquieran los documentos disponibles en el mercado editorial, sea directamente los documentos, o por medio de bases de datos científicas, pagando ellos mismos los costos asociados a dichos accesos.

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/282305791>

Mycotoxin Contamination of the Food Supply Chain – Implications for One Health Programme

Article in *Procedia Food Science* · December 2015

DOI: 10.1016/j.profoo.2015.09.053

CITATIONS

23

READS

328

3 authors:



Dragan Milićević

Institute of meat hygiene and technology Belgrade Serbia

94 PUBLICATIONS 1,104 CITATIONS

[SEE PROFILE](#)



Ksenija Nesic

Scientific Institute of Veterinary Medicine of Serbia

73 PUBLICATIONS 572 CITATIONS

[SEE PROFILE](#)



Sandra Jakšić

Scientific Veterinary Institute "Novi Sad"

103 PUBLICATIONS 603 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Network for Ecohealth and One Health - NEOH [View project](#)



451-03-9/2021-14/200050 [View project](#)



International 58th Meat Industry Conference “Meat Safety and Quality: Where it goes?”

Mycotoxin contamination of the food supply chain - Implications for One Health programme

Dragan Milicevic^{a,*}, Ksenija Nesic^b, Sandra Jaksic^c

^a*Institute of Meat Hygiene and Technology, Kacanskog 13, 11040 Belgrade, Serbia*

^b*Institute of Veterinary Medicine of Serbia, Auto put bb, 11070 Belgrade, Serbia*

^c*Scientific Veterinary Institute Novi Sad, Rumenicki put 20, 21000 Novi Sad, Serbia*

Abstract

Mycotoxins are a group of naturally occurring toxic chemical substances, produced by different fungal species, which can cause illness or even death due to their toxigenic, carcinogenic, mutagenic and teratogenic effects. Though more than 400 mycotoxins are known to have toxic effects on human when ingested along with contaminated food, mycotoxins like aflatoxins, ochratoxins, zeralenone, trichothecenes and fumonisins are the major mycotoxins influencing the public health and agriculture. Due to the global nature of the food supply and advances in analytical capabilities, mycotoxin contamination will continue to be an area of concern for regulatory agencies, the food industry, and consumers.

© 2015 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Peer-review under responsibility of scientific committee of The 58th International Meat Industry Conference (MeatCon2015)

Keywords: Mycotoxins; food supply chain; Public health

1. Mycotoxins in general

Mycotoxins are natural contaminants produced by a range of fungal species. Their common occurrence in food and feed poses a threat to the health of humans and animals. This threat is caused either by the direct contamination of agricultural commodities or by a “carry-over” of mycotoxins and their metabolites into animal tissues, milk, and

* Corresponding author. Tel.: +381-11-2650-655; fax: +381-11-2651-825

E-mail address: dragan@inmesbgd.com

eggs after feeding of contaminated feedingstuffs¹. The most important agro-economic and public health classes of mycotoxins are aflatoxins (AFT), ochratoxins (OTA), zeralenone (ZEA), trichothecenes, and fumonisins (FUMs) produced by species of *Fusarium*, *Penicillium* and *Aspergillus*. If these mycotoxins occur at a considerably high level in foods, they can cause toxic effects ranging from acute to chronic (mutagenic, teratogenic, carcinogenic) manifestations in humans and animals². The economic impact of mycotoxins include loss of human and animal life, increased health care and veterinary care costs, reduced livestock production, disposal of contaminated foods and feeds, and investment in research and applications to reduce severity of the mycotoxin problem³. Due to their deleterious effects on humans and farm animals, several countries have implemented regulations prescribing mycotoxin limits in several food commodities intended for consumption. In 1993, the WHO-International Agency for Research on Cancer^{4,5} evaluated the carcinogenic potential of AFT, OTA, TCT, ZEA, and FUMs.

1.1. Factor affecting production, contamination of foods and feeds by mycotoxins

Various factors operate interdependently to affect fungal colonization and/or production of the mycotoxins. D'Mello and MacDonald⁶ categorized the factors as physical factors (moisture, relative humidity, temperature and mechanical damage), chemical factors (carbon dioxide, oxygen, composition of substrate, pesticide and fungicides), and biological factors (plant variety, stress, insects, spore load). The biological factors have been further sub-categorized into intrinsic factors, including fungal species, strain specificity, strain variation, and instability of toxigenic properties. This is in part due to the multidisciplinary nature of mycotoxin research, which involves analytical chemistry, toxicology, taxonomic mycology, hygienic measures, microbial physiology, epidemiology, and increasingly weaponry. Some molds are capable of producing more than one mycotoxin and some mycotoxins are produced by more than one fungal species. Often more than one mycotoxin is found on a contaminated substrate. Therefore, the presence of potentially toxigenic fungi does not imply the presence of mycotoxins and vice versa⁸.

1.2. Occurrence and significance of mycotoxins in foods and feeds

Mycotoxicoses in humans or animals are characterized as food or feed related, non-contagious, non-transferable, non-infectious, and non-traceable to microorganisms other than fungi. Clinical symptoms usually subside upon removal of contaminated food or feed. Mycotoxin contamination is encountered in various environments. For instance, mycotoxins have been frequently identified in grain products, coffee beans, spices and nuts. In agriculture, mycotoxins can be present mainly in livestock feed. A wide range of commodities can be contaminated with mycotoxins both pre- and post-harvest. Serbia is largely a developing agriculture-based economy, so therefore, surveillance for mycotoxins and mycotoxigenic fungi is critical for maintaining high quality of feed (Table 1) and food (Table 2)^{9,10,11}.

1.2.1. Aflatoxins

AFs are difuranocoumarin derivatives produced by a polyketide pathway by many strains of *Aspergillus flavus*, *Aspergillus parasiticus* and the rare *Aspergillus nomius*, which contaminate agricultural commodities. They have toxic, carcinogenic, mutagenic and teratogenic effects in laboratory animals. The liver is the main target organ for AF toxicity and carcinogenicity. The rate of metabolism and products formed determine differences in species susceptibility to AF. In this instance, the presumed intermediate metabolite, AFTB1 8,9-epoxide forms an adduct with DNA and consequentially disrupts the transcriptional and translational processes. Approximately 95% of AFB₁ metabolites excreted in milk are in the form of AFM₁, though AFM₂, AFG₁ and AFB are also reported.

1.2.2. Ochratoxin

OTA is produced by fungi of the genera *Aspergillus* and *Penicillium*. The target organ for OTA is the kidneys and initial interest in this group of toxins was as a cause of porcine nephropathy. OTA has subsequently, been

associated with human disorders most noticeably Balkan Endemic Nephropathy in the former Yugoslavia, Chronic Interstitial Nephropathy and kidney tumours. Historically, consumption of pork has been a significant source of human exposure to ochratoxin A in these regions¹².

Table 1. Incidence of mycotoxins in investigated livestock feedingstuffs produced in Serbia during 2014.

Feed/ feedingstuffs	AFTB ₁		OTA		T-2/HT-2		ZEA		DON	
	N	n (%)	N	n (%)	N	n (%)	N	n (%)	N	n (%)
Poultry	189	73.0	27	29.6	38	39.5	14	28.6	32	53.1
Pigs	190	58.4	62	30.7	77	26.0	77	48.1	66	63.6
Cattle	103	59.2	4	0	5	20.0	9	77.8	5	40.0
Corn	100	37.0	39	20.5	48	18.8	52	53.9	50	76.0
Silage	14	71.4	3	0	-	-	8	100.0	12	66.7
Sunflower meal	8	15.0	1	0	-	-	2	0	5	0
Wheat	2	50.0	2	0	1	0	3	33.3	5	100.0

N - total number of analysed samples; n - percentage of positive samples.

Table 2. Occurrence of mycotoxins analysed in selected foodstuffs from Serbia.

Types of foodstuffs	AFTB ₁		AFTM ₁		DON		ZEA		FUM's		T-2		OTA	
	N	n	N	n	N	n	N	n	N	n	N	n	N	n
Milk	-	-	50	100	-	-	-	-	-	-	-	-	-	-
Cheese (white and hard type)	-	-	54	100	-	-	-	-	-	-	-	-	-	-
Wheat flour	15	0	-	-	15	86.7	15	33.3	15	0	15	26.7	15	0

N - total number of analysed samples; n - percentage of positive samples.

1.2.3. Zearalenone

ZEA is a phenolic resorcylic acid lactone with potent oestrogenic properties, produced primarily by *Fusarium*. ZEA is a non-steroidal oestrogen and its major metabolites α -zearalenol and β -zearalenol elicit significant oestrogenic activity in human and animals, corresponding to their binding affinities for hepatic, uterine, mammary and hypothalamic oestrogen receptors. Pigs are very sensitive to ZEA whereas poultry are very tolerant.

1.2.4. Fumonisin

A number of *Fusarium* species produce fumonisins, a group of some 12 compounds of which fumonisin B1 is the most studied and most toxic. The toxicity of FUMs largely reflects their ability to disrupt sphingolipid metabolism by inhibiting the enzyme ceramide synthase; an enzyme responsible for the acylation of sphinganine and sphingosine. It is now established that the FUMs cause equine leukoencephalomalacia and porcine pulmonary oedema, are carcinogenic in rats and may be a major aetiological factor in the incidence of human oesophageal cancer in southern Africa and China.

1.2.5. Trichothecenes

Several fungal genera are capable of producing TCT, however, most of them have been isolated from *Fusarium spp.* TCT comprise a vast group of over 100 fungal metabolites with the same basic structure. Examples of type A TCT include T-2 toxin (T-2) and HT-2 toxin (HT-2), and diacetoxyscirpenol (DAS). Fusarenone-X (FUX), deoxynivalenol (DON), and nivalenol (NIV) are some of the common naturally occurring type B TCT. At the

cellular level, the main toxic effect of TCT mycotoxins appears to be a primary inhibition of protein synthesis. TCT affect actively dividing cells such as those lining the gastrointestinal tract, the skin, lymphoid and erythroid cells.

2. Mycotoxin analysis

Mycotoxin analysis in food and feed is generally a multistep process comprised of sampling, sample preparation, toxin extraction from the matrix (usually with mixtures of water and polar organic solvents), extract clean-up and finally detection and quantitative determination. Proper sampling, milling and homogenisation procedures are pre-requisite for obtaining reliable analytical results because of the heterogeneous distribution of mycotoxins in food commodities.

3. Mycotoxin legislation and regulations

Legislation and regulation are constantly evolving issues. Many developing countries have realised that reducing mycotoxins levels in foods will not only reduce the financial burden on health care, but also confer international trade advantages such as exports to the attractive European markets. Factors fundamental to a country's ability to protect its population from mycotoxins include the political will to address mycotoxin exposure and capability to test food for contamination, which determines whether requirements can be enforced.

4. Prevention of mycotoxins

Management practices to maximize plant performance and decrease plant stress can decrease mycotoxin contamination substantially. This includes planting adapted varieties, proper fertilization, weed control, necessary irrigation, and proper crop rotation. For post-harvest mycotoxin control, prevention of conditions that favor fungal growth and subsequent toxin production needs to be considered, *i.e.* factors such as water activity of stored products, temperature, grain condition, gas composition of the intergranular air, microbial interactions, and presence of chemical or biological preservatives.

References

1. Milicevic D. Mycotoxins in the food chain – Old problems and new solution. *Tehn mesa* 2009;**50**(1-2): 99-111.
2. Richard JL. Some major mycotoxins and their mycotoxicoses—an overview. *Int J Food Microbiol* 2007;**119** (1-2):3-10.
3. Hussein HS, Jeffrey M. Brasel. Toxicity, metabolism, and impact of mycotoxins on humans and animals. *Toxicol* 2001;**167**:101–34.
4. IARC, 1993. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. Some Naturally Occurring Substances: Food Items and Constituents, Heterocyclic Aromatic Amines and Mycotoxins. IARC, Lyon, France.
5. IARC, 2002. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans Fumonisin B1. IARC, Lyon, France.
6. D'Mello JPF, MacDonald AMC. Mycotoxins. *Anim Feed Sci Technol* 1997;**69**:155-66.
7. Moss MO. The environmental factors controlling mycotoxin formation. In: Smith, J.E., Anderson, R.A. editors, *Mycotoxins and Animal Foods*. Boca Raton, Florida, USA: CRC Press; 1991. p 37–56.
8. Fung F, Clark RF. Health effects of mycotoxins: a toxicological overview. *J Toxicol Clin Toxicol* 2004;**42**(2):217-34.
9. Skrbic B, Zivancev J, Durisic-Mladenovic N, Godula M. Principal mycotoxins in wheat flour from the Serbian market: Levels and assessment of the exposure by wheat-based products. *Food Cont* 2012;**25**:389-96.
10. Skrbic B, Zivancev J, Antic I, Godula M. Levels of aflatoxin M1 in different types of milk collected in Serbia: Assessment of human and animal exposure. *Food Cont* 2014;**40**:113-19.
11. Skrbic B, Antic I, Zivancev J. Presence of aflatoxin M1 in white and hard cheese samples from Serbia. *Food Control* 2015;**(50)**: 111-17.
12. Milicevic D, Grubic M, Radicevic T, Stefanovic S, Jankovic S, Vranic V. Ochratoxin A residue in broiler tissues – risk assessment. *Tehn mesa* 2011;**52**(2):268–75.